

2004
Update



Peru

EPIDEMIOLOGICAL FACT SHEETS
ON HIV/AIDS AND SEXUALLY TRANSMITTED INFECTIONS



Joint United Nations Programme on HIV/AIDS

UNAIDS

UNHCR • UNICEF • WFP • UNDP • UNFPA
UNODC • ILO • UNESCO • WHO • WORLD BANK



**Pan American
Health
Organization**

Regional Office of the
World Health Organization

unicef 



World Health
Organization

HIV/AIDS estimates

In 2003 and during the first quarter of 2004, UNAIDS and WHO worked closely with national governments and research institutions to recalculate current estimates on people living with HIV/AIDS. These calculations are based on the previously published estimates for 1999 and 2001 and recent trends in HIV/AIDS surveillance in various populations. A methodology developed in collaboration with an international group of experts was used to calculate the new estimates on prevalence and incidence of HIV and AIDS deaths, as well as the number of children infected through mother-to-child transmission of HIV. Different approaches were used to estimate HIV prevalence in countries with low-level, concentrated or generalised epidemics. The current estimates do not claim to be an exact count of infections. Rather, they use a methodology that has thus far proved accurate in producing estimates that give a good indication of the magnitude of the epidemic in individual countries. However, these estimates are constantly being revised as countries improve their surveillance systems and collect more information.

Adults in this report are defined as women and men aged 15 to 49. This age range covers people in their most sexually active years. While the risk of HIV infection obviously continues beyond the age of 50, the vast majority of those who engage in substantial risk behaviours are likely to be infected by this age. The 15 to 49 range was used as the denominator in calculating adult HIV prevalence.

Estimated number of adults and children living with HIV/AIDS, end of 2003

These estimates include all people with HIV infection, whether or not they have developed symptoms of AIDS, alive at the end of 2003:

| | | | |
|---------------------|---------|----------------|-----|
| Adults and children | 82,000 | | |
| Low estimate | 40,000 | | |
| High estimate | 140,000 | | |
| Adults (15-49) | 80,000 | Adult rate (%) | 0.5 |
| Low estimate | 39,000 | Low estimate | 0.3 |
| High estimate | 130,000 | High estimate | 0.9 |
| Children (0-15) | | | |
| Low estimate | | | |
| High estimate | | | |
| Women (15-49) | 27,000 | | |
| Low estimate | 13,000 | | |
| High estimate | 44,000 | | |

Estimated number of deaths due to AIDS

Estimated number of adults and children who died of AIDS during 2003:

| | |
|----------------|-------|
| Deaths in 2003 | 4,200 |
| Low estimate | 2,100 |
| High estimate | 7,300 |

Estimated number of orphans

Estimated number of children who have lost their mother or father or both parents to AIDS and who were alive and under age 17 at the end of 2003:

| | |
|------------------------|--|
| Current living orphans | |
| Low estimate | |
| High estimate | |

Assessment of the epidemiological situation 2004

The main modes of HIV -the heterosexual and men who have sex with men (MSM)- account for 40% and 42% respectively. HIV infection is concentrated among the poor in coastal cities where two thirds of reported cases are in Lima and the adjoining Callao area.

The highest rates of infection have been reported in MSM, although the proportion of females is increasing. In 1998, a cohort study of close to 5,000 MSM detected HIV rates of 12.2% of those in Lima, 14.5% in Iquitos and 7.5% in Pucallpa (both cities in the Amazonian region), 2.7%-5.3% in coastal cities and 1.4% in the Andean city of Cuzco. Consistent condom use was reported by only 12%, and 46% of MSM also reported having sex with women. Among transvestites in this study, HIV prevalence was 35%.

Other high-risk populations had lower HIV prevalence. Nationwide, 0.6% to 2% of female sex workers are positive for HIV but among non-legal sex workers 10% were HIV-positive.

HIV rates in pregnant women are still well below 1% in Lima, and lower still in the provinces. Sentinel surveillance among 15-24-year-olds at maternity hospitals in Lima found 0.23%-0.58% HIV positive between 1996 and 1999. AIDS cases are increasing in 20-24-year-olds, indicating that HIV infection is occurring in teenagers

In 2003 a random two-stage cluster household sample of 18 to 29-year-old men and women was selected in each of 24 Peruvian cities with populations over 50,000. A consecutive sample of cohabiting sex partners of selected participants was also included. Consenting participants completed a face to face questionnaire and biological samples were collected for HIV and sexually transmitted diseases (STDs), testing venous blood for HIV and syphilis urine in men and self collected vaginal swabs in women to test for Chlamydia and N. gonorrhoeae. Oral fluid for HIV screening was obtained from participants not willing to provide blood samples, and urine was collected from women not willing to provide vaginal swabs.

A total of 15,259 selected persons agreed to participate, with an overall participation rate of 90.5% for the randomly selected participants. The participation rate for their sex partners was 88.1%. Blood or oral fluid was collected from 84% of the participants, urine was obtained from 87% of males, and vaginal swabs or urine were obtained from 84% of women.

Sex with other men was reported by 12% of men (ever), while 6% reported sex with other men within the last 12 months. Consistent condom use reported by men was 24% with casual partners, 45% with sex workers and 32% with other males. Consistent condom use reported by women was only 12% with casual partners.

HIV seroprevalence in men was 0.4%, with values ranging from 0 to 1.9% among the cities screened, and higher rates in cities located in the Amazon jungle in eastern Peru. Overall HIV seroprevalence in women was 0.1%, with a range from 0 to 0.4% among cities. No HIV infections could be detected using oral fluid.

The most important risk factor for HIV infection for males was sex with other men, and for females, sex in exchange for money. Prevalences for other STDs were 1%, 0.3%, and 4% for syphilis, gonorrhoea, and chlamydia respectively in males; and 0.9%, 0.8%, 6.8%, 5.2%, and 25% for syphilis, gonorrhoea, chlamydia, trichomoniasis, and bacterial vaginosis respectively for females.

UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance

Global Surveillance of HIV/AIDS and sexually transmitted infections (STIs) is a joint effort of WHO and UNAIDS. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, initiated in November 1996, guides respective activities. The primary objective of the Working Group is to strengthen national, regional and global structures and networks for improved monitoring and surveillance of HIV/AIDS and STIs. For this purpose, the Working Group collaborates closely with national AIDS programmes and a number of national and international experts and institutions. The goal of this collaboration is to compile the best information available and to improve the quality of data needed for informed decision-making and planning at national, regional, and global levels. The Epidemiological Fact Sheets are one of the products of this close and fruitful collaboration across the globe.

Within this framework, the Fact Sheets collate the most recent country-specific data on HIV/AIDS prevalence and incidence, together with information on behaviours (e.g. casual sex and condom use) which can spur or stem the transmission of HIV.

Not unexpectedly, information on all of the agreed upon indicators was not available for many countries in 2003. However, these updated Fact Sheets do contain a wealth of information which allows identification of strengths in currently existing programmes and comparisons between countries and regions. The Fact Sheets may also be instrumental in identifying potential partners when planning and implementing improved surveillance systems.

The fact sheets can be only as good as information made available to the UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance. Therefore, the Working Group would like to encourage all programme managers as well as national and international experts to communicate additional information to them whenever such information becomes available. The Working Group also welcomes any suggestions for additional indicators or information proven to be useful in national or international decision-making and planning.

Basic indicators

For consistency reasons the data used in the table below are taken from official UN publications.

| DEMOGRAPHIC DATA | YEAR | ESTIMATE | SOURCE |
|---|-----------|----------|---------------------------------|
| Total population (thousands) | 2004 | 27,567 | UN population division database |
| Female population aged 15-24 (thousands) | 2004 | 2,637 | UN population division database |
| Population aged 15-49 (thousands) | 2004 | 14,529 | UN population division database |
| Annual population growth rate (%) | 1992-2002 | 1.7 | UN population division database |
| % of population in urban areas | 2003 | 73.7 | UN population division database |
| Average annual growth rate of urban population | 2000-2005 | 2.0 | UN population division database |
| Crude birth rate (births per 1,000 pop.) | 2004 | 22.6 | UN population division database |
| Crude death rate (deaths per 1,000 pop.) | 2004 | 6 | UN population division database |
| Maternal mortality rate (per 100,000 live births) | 2000 | 410 | WHO (WHR2004)/UNICEF |
| Life expectancy at birth (years) | 2002 | 69.7 | World Health Report 2004, WHO |
| Total fertility rate | 2002 | 2.9 | World Health Report 2004, WHO |
| Infant mortality rate (per 1,000 live births) | 2000 | 32 | World Health Report 2004, WHO |
| Under 5 mortality rate (per 1,000 live births) | 2000 | 42 | World Health Report 2004, WHO |

| SOCIO-ECONOMIC DATA | YEAR | ESTIMATE | SOURCE |
|--|-----------|----------|-------------------------------|
| Gross national income, ppp, per capita (Int.\$) | 2002 | 4,800 | World Bank |
| Gross domestic product, per capita % growth | 2001-2002 | 3.7 | World Bank |
| Per capita total expenditure on health (Int.\$) | 2001 | 231 | World Health Report 2004, WHO |
| General government expenditure on health as % of total expenditure on health | 2001 | 55 | World Health Report 2004, WHO |
| Total adult illiteracy rate | 2000 | 10.1 | UNESCO |
| Adult male illiteracy rate | 2000 | 5.3 | UNESCO |
| Adult female illiteracy rate | 2000 | 14.8 | UNESCO |
| Gross primary school enrolment ratio, male | 2000/2001 | 128 | UNESCO |
| Gross primary school enrolment ratio, female | 2000/2001 | 126 | UNESCO |
| Gross secondary school enrolment ratio, male | 2000/2001 | 89 | UNESCO |
| Gross secondary school enrolment ratio, female | 2000/2001 | 82 | UNESCO |

Contact address

UNAIDS/WHO Working Group on Global
HIV/AIDS and STI Surveillance
20, Avenue Appia
CH - 1211 Geneva 27
Switzerland
Fax: +41-22-791-4834

email: hivstrategicinfo@who.int or
estimates@unaids.org

website: <http://www.who.int/hiv>
<http://www.unaids.org>

Extracts of the information contained in these fact sheets may be reviewed, reproduced or translated for research or private study but not for sale or for use in conjunction with commercial purposes. Any use of information in these fact sheets should be accompanied by the following acknowledgment "UNAIDS/WHO epidemiological fact sheets on HIV/AIDS and Sexually Transmitted Infections, 2004 Update".

HIV prevalence in different populations

This section contains information about HIV prevalence in different populations. The data reported in the tables below are mainly based on the HIV database maintained by the United States Bureau of the Census where data from different sources, including national reports, scientific publications and international conferences are compiled. To provide a simple overview of the current situation and trends over time, summary data are given by population group, geographical area (Major Urban Areas versus Outside Major Urban Areas), and year of survey. Studies conducted in the same year are aggregated and the median prevalence rates (in percentages) are given for each of the categories. The maximum and minimum prevalence rates observed, as well as the total number of surveys/sentinel sites, are provided with the median, to give an overview of the diversity of HIV-prevalence results in a given population within the country. Data by sentinel site or specific study from which the medians were calculated are printed at the end of this fact sheet.

The differentiation between the two geographical areas Major Urban Areas and Outside Major Urban Areas is not based on strict criteria, such as the number of inhabitants. For most countries, Major Urban Areas were considered to be the capital city and - where applicable - other metropolitan areas with similar socio-economic patterns. The term Outside Major Urban Areas considers that most sentinel sites are not located in strictly rural areas, even if they are located in somewhat rural districts.

HIV sentinel surveillance*


| Group | Area | | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | |
|---------------------------|---------------------------|-------------------|---------|------|------|-------|------|------|------|------|------|-------|-------|-------|-------|------|------|------|------|--|
| Pregnant women | Major urban areas | N-Sites | | | | | | | | | | 11.00 | 12.00 | 11.00 | 1.00 | | 1.00 | | | |
| | | Minimum | | | | | | | | | | | 0 | 0.27 | 0 | 0.30 | | 0.62 | | |
| | | Median | | | | | | | | | | | 0 | 0.62 | 0.18 | 0.30 | | 0.62 | | |
| | | Maximum | | | | | | | | | | | 0.57 | 1.05 | 0.51 | 0.30 | | 0.62 | | |
| | Outside major urban areas | N-Sites | | | | | | | | | | | 5.00 | 13.00 | 12.00 | | | | | |
| | | Minimum | | | | | | | | | | | 0 | 0 | 0 | | | | | |
| | | Median | | | | | | | | | | | 0 | 0 | 0 | | | | | |
| | | Maximum | | | | | | | | | | | 0.17 | 0.98 | 0.51 | | | | | |
| | Sex workers | Major urban areas | N-Sites | 1.00 | | | 1.00 | | | | | | 1.00 | | 1.00 | | | | | |
| Minimum | | | 0.73 | | | 0.29 | | | | | | | 5.00 | | 1.60 | | | | | |
| Median | | | 0.73 | | | 0.29 | | | | | | | 5.00 | | 1.60 | | | | | |
| Maximum | | | 0.73 | | | 0.29 | | | | | | | 5.00 | | 1.60 | | | | | |
| Outside major urban areas | | N-Sites | 1.00 | 1.00 | 1.00 | | | 1.00 | | 1.00 | | 1.00 | 1.00 | 1.00 | | | 1.00 | | | |
| | | Minimum | 0.31 | 0.26 | 0.63 | | | 0.60 | | 0.80 | | 0.80 | 1.85 | 1.60 | | | 1.20 | | | |
| | | Median | 0.31 | 0.26 | 0.63 | | | 0.60 | | 0.80 | | 0.80 | 1.85 | 1.60 | | | 1.20 | | | |
| | | Maximum | 0.31 | 0.26 | 0.63 | | | 0.60 | | 0.80 | | 0.80 | 1.85 | 1.60 | | | 1.20 | | | |
| Injecting drug users | | Major urban areas | N-Sites | 1.00 | | | 1.00 | | | | | | | | | | | | | |
| | Minimum | | 1.23 | | | 28.13 | | | | | | | | | | | | | | |
| | Median | | 1.23 | | | 28.13 | | | | | | | | | | | | | | |
| | Maximum | | 1.23 | | | 28.13 | | | | | | | | | | | | | | |
| STI patients | Major urban areas | N-Sites | 1.00 | | | 1.00 | | | | | | | | | | | | | | |
| | | Minimum | 1.74 | | | 18.68 | | | | | | | | | | | | | | |
| | | Median | 1.74 | | | 18.68 | | | | | | | | | | | | | | |
| | | Maximum | 1.74 | | | 18.68 | | | | | | | | | | | | | | |
| | Outside major urban areas | N-Sites | | | | | | | | | | 1.00 | | | | | | | | |
| | | Minimum | | | | | | | | | | 7.00 | | | | | | | | |
| | | Median | | | | | | | | | | 7.00 | | | | | | | | |
| | | Maximum | | | | | | | | | | 7.00 | | | | | | | | |
| | Men having sex with men | Major urban areas | N-Sites | 1.00 | | | 1.00 | | | | | | 1.00 | 1.00 | | 1.00 | | | | |
| Minimum | | | 7.96 | | | 41.37 | | | | | | 18.20 | 20.66 | | 12.20 | | | | | |
| Median | | | 7.96 | | | 41.37 | | | | | | 18.20 | 20.66 | | 12.20 | | | | | |
| Maximum | | | 7.96 | | | 41.37 | | | | | | 18.20 | 20.66 | | 12.20 | | | | | |
| Outside major urban areas | | N-Sites | | | | | | | | | | | | 1.00 | 6.00 | | | | | |
| | | Minimum | | | | | | | | | | | | 1.15 | 1.40 | | | | | |
| | | Median | | | | | | | | | | | | 1.15 | 4.35 | | | | | |
| | | Maximum | | | | | | | | | | | | 1.15 | 14.50 | | | | | |
| Tuberculosis patients | | | | | | | | | | | | | | | | | | | | |

*Detailed data by site can be found in the Annex.



Maps & charts

Mapping the geographical distribution of HIV prevalence among different population groups may assist in interpreting both the national coverage of the HIV surveillance system as well in explaining differences in levels of prevalence. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, in collaboration with the WHO Public Health Mapping Team, Communicable Diseases, is producing maps showing the location and HIV prevalence in relation to population density, major urban areas and communication routes. For generalized epidemics, these maps show the location of prevalence of antenatal surveillance sites.

Trends in antenatal sentinel surveillance for higher prevalence countries, or in prevalence among selected populations for countries with concentrated epidemics, are a new addition. These are presented for those countries where sufficient data exist.



Trends in HIV prevalence in high risk groups



Median prevalence and ranges are shown in areas with more than one sentinel site.

The boundaries and names shown and the designations used on the map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement. WHO 2004, all rights reserved.

Reported AIDS cases

Following WHO and UNAIDS recommendations, AIDS case reporting is carried out in most countries. Data from individual AIDS cases are aggregated at the national level and sent to WHO. However, case reports come from surveillance systems of varying quality. Reporting rates vary substantially from country to country and low reporting rates are common in developing countries due to weaknesses in the health care and epidemiological systems. In addition, countries use different AIDS case definitions. A main disadvantage of AIDS case reporting is that it only provides information on transmission patterns and levels of infection approximately 5-10 years in the past, limiting its usefulness for monitoring recent HIV infections.

Despite these caveats, AIDS case reporting remains an important advocacy tool and is useful in estimating the burden of HIV-related morbidity as well as for short-term planning of health care services. AIDS case reports also provide information on the demographic and geographic characteristics of the affected population and on the relative importance of the various exposure risks. In some situations, AIDS reports can be used to estimate earlier HIV infection patterns using back-calculation. AIDS case reports and AIDS deaths have been dramatically reduced in industrialized countries with the introduction of Anti-Retroviral Therapy (ART).

| | | | | | | | | | | | | | | | | | | | |
|------|------|------|------|------|-------|------|---------------------|------|------|------|------|------|------|------|------|------|------|------|------|
| 1979 | 1980 | 1981 | 1982 | 1983 | 1984 | 1985 | 1986 | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 |
| | | | | 1 | 4 | 4 | 22 | 97 | 250 | 317 | 410 | 480 | 748 | 709 | 840 | 1090 | 1177 | 1078 | 1031 |
| 1999 | 2000 | 2001 | 2002 | 2003 | Total | UNK | Date of last report | | | | | | | | | | | | |
| 1009 | 615 | | | | 9882 | | 11/11/2001 | | | | | | | | | | | | |

Curable sexually transmitted infections (STIs)

The predominant mode of transmission of both HIV and other STIs is sexual intercourse. Measures for preventing sexual transmission of HIV and STIs are the same, as are the target audiences for interventions. In addition, strong evidence supports several biological mechanisms through which STIs facilitate HIV transmission by increasing both HIV infectiousness and HIV susceptibility. Thus, detection and treatment of individuals with STIs is an important part of an HIV control strategy. In summary, if the incidence/prevalence of STIs is high in a country, then there is the possibility of high rates of sexual transmission of HIV. Monitoring trends in STIs provides valuable insight into the likelihood of the importance of sexual transmission of HIV within a country, and is part of second generation surveillance. These trends also assist in assessing the impact of behavioural interventions, such as delaying sexual debut, reducing the number of sex partners and promoting condom use.

Clinical services offering STI care are an important access point for people at high risk for both STIs and HIV. Identifying people with STIs allows for not only the benefit of treating the STI, but for prevention education, HIV testing, identifying HIV-infected persons in need of care, and partner notification for STIs or HIV infection. Consequently, monitoring different components of STI prevention and control can also provide information on HIV prevention and control activities within a country.

STI syndromes

| Reported cases | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | Incidence 2003 |
|----------------|------|------|------|------|------|------|------|------|----------------|
|----------------|------|------|------|------|------|------|------|------|----------------|

Comments:

Source:

Syphilis prevalence, women

Percent of blood samples taken from pregnant women aged 15-49 that test positive for syphilis - positive reaginic and treponemal test-during routine screening at selected antenatal clinics.

| Year | Area | Rate | Range |
|------|------|------|-------|
|------|------|------|-------|

Comments:

Source:

Estimated prevalence of curable STIs among female sex workers

- Chlamydia

| Year | Area | Rate | Range |
|-----------|-------|------|-------|
| 1997-1999 | Urban | 22 | |

Comments:

Source: Paris M. Prevalence of gonococcal and chlamydial infections in commercial sex workers in a Peruvian Amazon city. Sex Transm Dis 1999; 26(2):103-107.

- Gonorrhoea

| Year | Area | Rate | Range |
|-----------|-------|------|-------|
| 1997-1999 | Urban | 14 | |

Comments:

Source: Paris M. Prevalence of gonococcal and chlamydial infections in commercial sex workers in a Peruvian Amazon city. Sex Transm Dis 1999; 26(2):103-107.

Estimated prevalence of curable STIs among female sex workers (continued)**- Syphilis**

| Year | Area | Rate | Range |
|-------------|---------------|-------------|--------------|
| 2000-2001 | Not specified | 6.5 | 0.1-23.4 |

Comments:

Source: Sexually Transmitted Diseases Prevalences from the 2002 National Household-Based General Population Survey in Young Urban Adults in Peru.

- Trichomoniasis

| Year | Area | Rate | Range |
|-------------|-------------|-------------|--------------|
|-------------|-------------|-------------|--------------|

Comments:

Source:

Health service and care indicators

HIV prevention strategies depend on the twin efforts of care and support for those living with HIV or AIDS, and targeted prevention for all people at risk or vulnerable to the infection. It is difficult to capture such a large range of activities with one or just a few indicators. However, a set of well-established health care indicators may help to identify general strengths and weaknesses of health systems. Specific indicators, such as access to testing and blood screening for HIV, help to measure the capacity of health services to respond to HIV/AIDS - related issues.

Access to health care

| Indicators | Year | Estimate | Source |
|--|------|----------|--------------|
| % of population with access to health services - total | | | |
| % of population with access to health services - urban | | | |
| % of population with access to health services - rural | | | |
| Contraceptive prevalence rate (%) | 1996 | 64.2 | UNICEF/UNPOP |
| Percentage of contraceptive users using condoms | | | |
| % of births attended by skilled health personnel | 2000 | 56.4 | WHO |
| % of 1-yr-old children fully immunized - DPT | 2002 | 89 | WHO/UNICEF |
| % of 1-yr-old children fully immunized - Measles | 2001 | 97 | WHO/UNICEF |
| % of ANC clinics where HIV testing is available | | | |

Number of adults (15-49) with advanced HIV infection receiving ARV therapy as of June 2004

Adults on treatment

Number: 1,900

Source: WHO

Estimated number of adults (15-49) in need of treatment in 2003

Adults needing treatment

Number: 9,700

Source: WHO/UNAIDS

Coverage of HIV testing and counselling

Number of public and NGO services providing testing and counselling services.

| Year | Area | N= |
|------|------|----|
|------|------|----|

Comments:

Source:

Knowledge and behaviour

In most countries the HIV epidemic is driven by behaviours (e.g.: multiple sexual partners, injecting drug use) that expose individuals to the risk of infection. Information on knowledge and on the level and intensity of risk behaviour related to HIV/AIDS is essential in identifying populations most at risk for HIV infection and in better understanding the dynamics of the epidemic. It is also critical information in assessing changes over time as a result of prevention efforts. One of the main goals of the 2nd generation HIV surveillance systems is the promotion of a standard set of indicators defined in the National Guide (Source: National AIDS Programmes, A Guide to Monitoring and Evaluation, UNAIDS/00.17) and regular behavioural surveys in order to monitor trends in behaviours and to target interventions.

The indicators on knowledge and misconceptions are an important prerequisite for prevention programmes to focus on increasing people's knowledge about sexual transmission, and, to overcome the misconceptions that act as a disincentive to behaviour change. Indicators on sexual behaviour and the promotion of safer sexual behaviour are at the core of AIDS programmes, particularly with young people who are not yet sexually active or are embarking on their sexual lives, and who are more amenable to behavioural change than adults. Finally, higher risk male-male sex reports on unprotected anal intercourse, the highest risk behaviour for HIV among men who have sex with men.

Knowledge of HIV prevention methods

Prevention indicator: Percentage of young people 15-24 who both correctly identify two ways of preventing the sexual transmission of HIV and who reject three misconceptions about HIV transmission.

| Year | Male | Female |
|------|------|--------|
| | | |

Comments:

Source:

Reported condom use at last higher risk sex (young people 15-24)

Prevention indicator: Proportion of young people reporting the use of a condom during sex with a non-regular partner.

| Year | Male | Female |
|------|------|--------|
| 2000 | | 19 |

Comments:

Source: DHS

Age-mixing in sexual partnerships among young women

The proportion of young women who have had sex in the last 12 months with a partner who is 10 or more years older than themselves.

| Year | Area | Age group | Male | Female | All |
|------|------|-----------|------|--------|-----|
| | | | | | |

Comments:

Source:

Reported non-regular sexual partnerships

Prevention indicator: Proportion of young people 15-24 having at least one sex partner other than a regular partner in the last 12 months.

| Year | Male | Female |
|------|------|--------|
| 2000 | | 29 |

Comments:

Source: DHS

Knowledge and behaviour (continued)Ever used a condom

Percentage of people who ever used a condom.

| Year | Area | Age group | Male | Female | All |
|------|------|-----------|------|--------|-----|
|------|------|-----------|------|--------|-----|

Comments:

Source:

Adolescent pregnancy

Percentage of teenagers 15-19 who are mothers or pregnant with their first child.

| Year | Percentage |
|------|------------|
|------|------------|

Comments:

Source:

Age at first sexual experience

Proportion of 15-19 year olds who have had sex before age 15.

| Year | Male | Female |
|------|------|--------|
| 2000 | | 5 |

Comments:

Source: DHS

Prevention indicators

Male and female condoms are the only technology available that can prevent sexual transmission of HIV and other STIs. Persons exposing themselves to the risk of sexual transmission of HIV should have consistent access to high quality condoms. AIDS Programs implement activities to increase both availability of and access to condoms. These activities should be monitored and have resources directed to problem areas. The indicator below highlights the availability of condoms. However, even if condoms are widely available, this does not mean that individuals can or do access them.

Condom availability nationwide

Total number of condoms available for distribution nationwide during the preceding 12 months, divided by the total population aged 15-49.

| Year | N | Rate |
|------|---|------|
|------|---|------|

Comments:

Source:

Prevention of mother-to-child transmission (MTCT) nationwide

Percentage of women who were counselled during antenatal care for their most recent pregnancy, accepted an offer of testing and received their test results, of all women who were pregnant at any time in the preceding two years.

| Year | N | Rate |
|------|---|------|
|------|---|------|

Comments:

Source:

Blood safety programs aim to ensure that the majority of blood units are screened for HIV and other infectious agents. This indicator gives an idea of the overall percentage of blood units that have been screened to high enough standards that they can confidently be declared free of HIV.

Screening of blood transfusions nationwide

Percentage of blood units transfused in the last 12 months that have been adequately screened for HIV according to national or WHO guidelines.

| Year | N | Rate |
|------|---|------|
|------|---|------|

Comments:

Source:

Sources

Data presented in this Epidemiological Fact Sheet come from several sources, including global, regional and country reports, published documents and articles, posters and presentations at international conferences, and estimates produced by UNAIDS, WHO and other United Nations agencies. This section contains a list of the more relevant sources used for the preparation of the Fact Sheet. Where available, it also lists selected national Web sites where additional information on HIV/AIDS and STI are presented and regularly updated. However, UNAIDS and WHO do not warrant that the information in these sites is complete and correct and shall not be liable whatsoever for any damages incurred as a result of their use.

Aguero, G., F. S. Wignall, W. Alexander, et al. 1988 HIV Infections in Peru IV International Conference on AIDS, Stockholm, 6/13-14, Poster 5078.

Alarcon, J., O. Palacios, A. Tejada, et al. 1989 Incidence d'Infection HIV-1 dans Prostituees de Callao, Perou. Etude de Cohorte. V International Conference on AIDS, Montreal, 6/4-9, Poster Th.H.P. 17.

Alarcon, J., C. Rodriguez, K. Johnson, et al. 1997 Determinants and Prevalence of HIV and Syphilis in Pregnant Women in Lima, Peru V Pan-American Conference on AIDS and XI Latin American Congress on STD, Lima, Peru, 12/3-6, Abstract P033.

Carcamo, C., P. Minaya, M. Chiong, et al. 2000 HIV Sentinel Surveillance in Peru XIII International AIDS Conference, Durban, South Africa, 7/9-14, Abstract ThOrC677.

Golenbock, D. T., J. Guerra, J. Pfister, et al. 1988 Absence of Infection with Human Immunodeficiency Virus in Peruvian Prostitutes AIDS Research and Human Retroviruses, vol. 4, no. 6, pp. 493-499.

Getaneh, A., J. Sanchez, D. Watts, et al. 1997 Sero Prevalence of Sexually Transmitted Disease and Infections Related to Sexual Behavioral Characteristics among the ... V Pan-American Conference on AIDS and XI Latin American Congress on STD, Lima, Peru, 12/3-6, Abstract PCV264.

Jorge, S., K. Russell, C. Carcamo, et al. 2000 HIV Sentinel Surveillance for Men Who have Sex with Men in Peru XIII International AIDS Conference, Durban, South Africa, 7/9-14, Abstract ThOrC717.

Lama, J., P. Goicochea, M. Chion, et al. 1997 Comportamiento Sexual y Factores de Riesgo Para ETS e Infeccion VIH Entre Hombres que Tienen Sexo Con Otros ... V Pan-American Conference on AIDS and XI Latin American Congress on STD, Lima, Peru, 12/3-6, Abstract P086.

McCarthy, M. C., F. S. Wignall, J. Sanchez, et al. 1996 The Epidemiology of HIV-1 Infection in Peru, 1986-1990 AIDS vol. 10, no. 10, pp. 1141-1145.

Melvin, A. J., J. Alarcon, C. Velasquez, et al. 2002 Rapid HIV-1 Testing of Pregnant Women Presenting in Labor with Unknown HIV Status in Lima, Peru 9th Conference on Retroviruses and Opportunistic Infections, Foundation for Retrovirology and Human Health, Seattle, WA., 2/24-28, online abstract, Poster Session 104, #789-W.

Peru Ministerio de Salud 1999 Informacion Estadistica: 1983-Julio-1999 Programa de Control de Enfermedades de Transmision Sexual Y Sida (PROCETSS), report.

Peru 1999 Pregnant Women Sentinel Surveillance - Peru 1996 - 1999 Personal communication, unpublished tables.

Quijano, E., M. Cairampoma, A. Ramirez, et al. 1997 Impacto de las Intervenciones de Salud Sobre la Infeccion por VIH en Trabajadoras Sexuales Callao, Peru 1987 - Junio 1997 V Pan-American Conference on AIDS and XI Latin American Congress on STD, Lima, Peru, 12/3-6, Abstract PCJ176.

Rojas, G., E. Gotuzzo, A. Yi, et al. 1986 Acquired Immunodeficiency Syndrome in Peru Annals of Internal Medicine, vol. 105, no. 3, pp. 465.

Revata, C., P. Quispe, E. Nino de Guzman, et al. 1997 Seroprevalencia de Infeccion por VIH en 2 Grupos de Frecuente Transmision en la Provincia de ICA, Peru 1995 - 1996 V Pan-American Conference on AIDS and XI Latin American Congress on STD, Lima, Peru, 12/3-6, Abstract PCV256.

Sanchez, J. 2000 Pregnant Women Sentinel Surveillance - Peru 1996-1999 UNAIDS Epifact Sheet Update.

Suarez-Ognio, L. A. N., H. Manrique-Chicama, M. M. Pun-Chinarro 2002 Hypothetical Model of Transmission Dynamic of HIV/AIDS in Peru XIV International AIDS Conference, Barcelona, Spain, 7/7-12, Abstract WePeC6108.

Tabet, S., J. Sanchez, J. Lama, et al. 2002 HIV, Syphilis and Heterosexual Bridging among Peruvian Men who Have Sex with Men AIDS, vol. 16, no. 9, pp. 1271-1277.

Websites: **National AIDS Programme (Spanish only):**
<http://www.minsa.gob.pe/procetss/index.htm>

Annex: HIV surveillance by site

| Group | Area | 1987 | 1988 | 1989 | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | | |
|-----------------------|---------------------------|-----------------------------------|---------------------------|----------|------|------|-------|-------|------|------|------|-------|-------|-------|-------|------|------|------|------|--|
| Pregnant women | Major urban areas | Honodami San Bartolome, Lima | | | | | | | | | 0.33 | 0.33 | 0.33 | | | | | | | |
| | | Hospital Arzobispo Loayza, Lima | | | | | | | | | | 0.55 | 0.98 | 0.30 | | | | | | |
| | | Hospital Cayetano Heredia, Lima | | | | | | | | | | 0.57 | 0.66 | 0.51 | | | | | | |
| | | Hospital Daniel A Carrion, Lima | | | | | | | | | | 0 | 0.31 | 0 | | | | | | |
| | | Hospital Dos de Mayo, Lima | | | | | | | | | | 0 | 0.66 | 0 | | | | | | |
| | | Hospital Hipolito Unanue, Lima | | | | | | | | | | | 0 | 0.30 | 0 | | | | | |
| | | Hospital Maria Auxiliadora, Lima | | | | | | | | | | | 0 | 0.27 | 0.18 | | | | | |
| | | Hospital Santa Rosa, Lima | | | | | | | | | | | 0 | 1.05 | 0 | | | | | |
| | | Hospital Sergio Bernales, Lima | | | | | | | | | | | 0 | 0.71 | 0 | | | | | |
| | | Instituto Materno-Perinatal, Lima | | | | | | | | | | | | 0.47 | | | | | 0.62 | |
| | Lima | | | | | | | | | | | 0.28 | 0.58 | 0.23 | 0.30 | | | | | |
| | Maternidad de Lima, Lima | | | | | | | | | | | 0.52 | 0.68 | 0.34 | | | | | | |
| | Outside major urban areas | Arequipa | | | | | | | | | | 0 | 0 | 0 | | | | | | |
| | | Chiclayo | | | | | | | | | | 0.17 | 0 | 0 | | | | | | |
| | | Chimbote | | | | | | | | | | | 0 | 0 | | | | | | |
| | | Cusco | | | | | | | | | | 0 | 0 | 0 | | | | | | |
| | | Huancayo | | | | | | | | | | | | 0.98 | 0 | | | | | |
| | | Ica | | | | | | | | | | | 0 | 0.51 | | | | | | |
| | | Iquitos | | | | | | | | | | 0 | 0.32 | 0 | | | | | | |
| | | Piura | | | | | | | | | | | | 0 | 0 | | | | | |
| Sullana | | | | | | | | | | | | | 0 | 0 | | | | | | |
| Tacna | | | | | | | | | | | | | 0.46 | 0 | | | | | | |
| Sex workers | Major urban areas | Lima | | | | | | | | | 5.00 | | 1.60 | | | | | | | |
| | | National | 0.73 | | | 0.29 | | | | | | | | | | | | | | |
| | Outside major urban areas | Callao | 0.31 | | 0.63 | | | | | | | | | | | | | | | |
| | | Ica | | | | | | | | | | | | 1.85 | | | | | | |
| | | Not specified | | 0.26 | | | | 0.60 | 0.80 | | | 0.80 | | 1.60 | | 1.20 | | | | |
| | | Major urban areas | National | 1.23 | | | 28.13 | | | | | | | | | | | | | |
| | | Outside major urban areas | Not specified | | | | | | | 7.00 | | | | | | | | | | |
| | | Men having sex with men | Major urban areas | Lima | | | | | | | | 18.20 | 20.66 | | 12.20 | | | | | |
| | | | | National | 7.96 | | | 41.37 | | | | | | | | | | | | |
| | | | Outside major urban areas | Chiclayo | | | | | | | | | | | | 2.70 | | | | |
| Cusco | | | | | | | | | | | | | | 1.40 | | | | | | |
| Ica | | | | | | | | | | | | | 1.15 | | | | | | | |
| Iquitos | | | | | | | | | | | | | | 14.50 | | | | | | |
| Pucallpa | | | | | | | | | | | | | | 7.50 | | | | | | |
| Sullana | | | | | | | | | | | | | | 5.30 | | | | | | |
| Tacna | | | | | | | | | | | | | | 3.40 | | | | | | |
| Tuberculosis patients | | | | | | | | | | | | | | | | | | | | |