

NPBI-8

Young people's condom use with non-regular partners*

Data source: name

Data source: type

Data collection period (day/month/year) to

PART I: Data requirements	Males			Females			Both sexes		
	Urban	Rural	National	Urban	Rural	National	Urban	Rural	National

NUMERATOR

Instructions:

- i) Calculate median age at first sex (UNAIDS, *National AIDS Programmes: A Guide to Monitoring and Evaluation*); enter in Line 1.
- ii) Select *only* those respondents (aged 15-24) who gave answers (excluding "don't know") to *all* of questions 2 to 5 below
- iii) Line 2: enter the number of respondents who stated that they had commenced sexual activity
- iv) Line 3: enter the number of respondents who stated that they had had any form of sexual relationship in the last 12 months
- v) Line 4: enter the number of respondents who stated that they had had a non-regular sexual partner in the last 12 months (NB: a 'non-regular' sexual partner here is someone the respondent was not married to and not cohabiting with at the time they had sex)
- vi) Line 5: enter the number of respondents who answered "yes" to the question in line 2 and who reported using condom when they last had sex with this non-regular partner

1. Median age at first sex*	<input type="text"/>	<input type="text"/>	NA	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>		
2. Commenced sexual activity	<input type="text"/>	<input type="text"/>		<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>		
3. Sexual partner within the last 12 months*	<input type="text"/>	<input type="text"/>		<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>		
4. Non-regular sexual partner within the last 12 months*	<input type="text"/>	<input type="text"/>	3,3%	<input type="text"/>	<input type="text"/>	0,6%	<input type="text"/>	<input type="text"/>	3,9%
5. Had a non-regular sexual partner within the last 12 months and used a condom the last time had sex with this partner Data from KABP study/ Sample 3000	<input type="text"/>	<input type="text"/>	40,3%	<input type="text"/>	<input type="text"/>	NA	<input type="text"/>	<input type="text"/>	

DENOMINATOR

6. Numbers of respondents (aged 15-24) who reported having had a non-regular sexual partner in the last 12 months (i.e., line 4 above)	<input type="text"/>	<input type="text"/>		<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
7. Percentage of the national population (aged 15-24) who live in urban areas**			<input type="text"/>			<input type="text"/>	<input type="text"/>

PART II:

Indicator computation

INDICATOR SCORES BY SEX & RESIDENCE

8. Divide the number of respondents who reported using condoms with their last non-regular partner (line 5) by the number who reported having had a non-regular sexual partner in the last 12 months (line 6) and multiply the result by 100.	<input type="text" value="####"/>	<input type="text" value="####"/>	<input type="text" value="####"/>	<input type="text" value="####"/>	<input type="text" value="####"/>	<input type="text" value="####"/>
---	-----------------------------------	-----------------------------------	-----------------------------------	-----------------------------------	-----------------------------------	-----------------------------------

INDICATOR SCORES BY SEX (NATIONAL)

9. i) Calculate the weighted average of the urban and rural indicator scores (line 8) using the percentages who live in urban and rural areas (line 7) as the weights. ii) Take the simple average of the national scores for men and women to get the combined score.	<input type="text" value="####"/>	<input type="text" value="####"/>	<input type="text" value="####"/>
---	-----------------------------------	-----------------------------------	-----------------------------------

* The data collected here also provide the information needed for the first two additional indicators recommended in the UNGASS indicator. guidelines

** From National Census Office statistics (as for NPI-1).