

2004  
Update



## Philippines

**EPIDEMIOLOGICAL FACT SHEETS**  
ON HIV/AIDS AND SEXUALLY TRANSMITTED INFECTIONS



Joint United Nations Programme on HIV/AIDS

**UNAIDS**

UNHCR • UNICEF • WFP • UNDP • UNFPA  
UNODC • ILO • UNESCO • WHO • WORLD BANK



World Health  
Organization

## HIV/AIDS estimates

In 2003 and during the first quarter of 2004, UNAIDS and WHO worked closely with national governments and research institutions to recalculate current estimates on people living with HIV/AIDS. These calculations are based on the previously published estimates for 1999 and 2001 and recent trends in HIV/AIDS surveillance in various populations. A methodology developed in collaboration with an international group of experts was used to calculate the new estimates on prevalence and incidence of HIV and AIDS deaths, as well as the number of children infected through mother-to-child transmission of HIV. Different approaches were used to estimate HIV prevalence in countries with low-level, concentrated or generalised epidemics. The current estimates do not claim to be an exact count of infections. Rather, they use a methodology that has thus far proved accurate in producing estimates that give a good indication of the magnitude of the epidemic in individual countries. However, these estimates are constantly being revised as countries improve their surveillance systems and collect more information.

Adults in this report are defined as women and men aged 15 to 49. This age range covers people in their most sexually active years. While the risk of HIV infection obviously continues beyond the age of 50, the vast majority of those who engage in substantial risk behaviours are likely to be infected by this age. The 15 to 49 range was used as the denominator in calculating adult HIV prevalence.

### Estimated number of adults and children living with HIV/AIDS, end of 2003

These estimates include all people with HIV infection, whether or not they have developed symptoms of AIDS, alive at the end of 2003:

Adults and children	9,000		
Low estimate	3,000		
High estimate	18,000		
Adults (15-49)	8,900	Adult rate (%)	<0.1
Low estimate	2,900	Low estimate	
High estimate	18,000	High estimate	<0.2
Children (0-15)			
Low estimate			
High estimate			
Women (15-49)	2,000		
Low estimate	700		
High estimate	4,000		

### Estimated number of deaths due to AIDS

Estimated number of adults and children who died of AIDS during 2003:

Deaths in 2003	<500
Low estimate	
High estimate	<1,000

### Estimated number of orphans

Estimated number of children who have lost their mother or father or both parents to AIDS and who were alive and under age 17 at the end of 2003:

Current living orphans
Low estimate
High estimate

## Assessment of the epidemiological situation 2004

The Philippines remains a low HIV prevalence country. Since the first cases of HIV/AIDS were reported in 1984, 1,515 HIV infections, including 508 AIDS cases and 196 HIV/AIDS-related deaths, had been reported as of June 2001. No significant epidemic HIV transmission has yet been detected in the Philippines and, as a result, the official national HIV prevalence estimate has been reduced from an initial 50 000 in the early 1990s to 26 000 a few years ago and, most recently, in 2003, to about 9 000.

HIV in the Philippines is predominantly sexually transmitted (90%). The number of HIV/AIDS cases is not expected to increase substantially over the next few years.

There is a high prevalence of STI among sex workers, with the prevalence rate of selected STIs--predominantly chlamydial infections--reaching more than 40%. The prevalence rate of STIs among women attending antenatal clinics was <1% for STI other than chlamydia (5.6%). Gonococcal resistance to penicillin, tetracycline, and ciprofloxacin is high. Most of gonorrhoea isolates (89.3%) were resistant to penicillin and 38% were resistant to quinolones in 2000.

The first large-scale behavioural sentinel surveillance (BSS) was conducted in 1997. There has been an increasing trend towards condom use during the last sex with a non-regular partner among registered sex workers from 75% in 1997 to 76% in 2000 and 83% in 2002. As a result of outreach interventions, the rate among freelance sex workers increased from 65% in 1997 to 72% in 2000 and 80% in 2002. Sex workers have, on average, a low number of clients (2-4 per week in 2002). A recent study indicated that 13% of men had two or more sex partners. There was a statistically significant decline in the proportion of IDU in Cebu who shared injecting equipment from 77% in 1998 to 53% in 2000, but this decline was reversed in 2001 when sharing increased to 81%.

## UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance

Global Surveillance of HIV/AIDS and sexually transmitted infections (STIs) is a joint effort of WHO and UNAIDS. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, initiated in November 1996, guides respective activities. The primary objective of the Working Group is to strengthen national, regional and global structures and networks for improved monitoring and surveillance of HIV/AIDS and STIs. For this purpose, the Working Group collaborates closely with national AIDS programmes and a number of national and international experts and institutions. The goal of this collaboration is to compile the best information available and to improve the quality of data needed for informed decision-making and planning at national, regional, and global levels. The Epidemiological Fact Sheets are one of the products of this close and fruitful collaboration across the globe.

Within this framework, the Fact Sheets collate the most recent country-specific data on HIV/AIDS prevalence and incidence, together with information on behaviours (e.g. casual sex and condom use) which can spur or stem the transmission of HIV.

Not unexpectedly, information on all of the agreed upon indicators was not available for many countries in 2003. However, these updated Fact Sheets do contain a wealth of information which allows identification of strengths in currently existing programmes and comparisons between countries and regions. The Fact Sheets may also be instrumental in identifying potential partners when planning and implementing improved surveillance systems.

The fact sheets can be only as good as information made available to the UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance. Therefore, the Working Group would like to encourage all programme managers as well as national and international experts to communicate additional information to them whenever such information becomes available. The Working Group also welcomes any suggestions for additional indicators or information proven to be useful in national or international decision-making and planning.

## Basic indicators

For consistency reasons the data used in the table below are taken from official UN publications.

DEMOGRAPHIC DATA	YEAR	ESTIMATE	SOURCE
Total population (thousands)	2004	81,408	UN population division database
Female population aged 15-24 (thousands)	2004	8,130	UN population division database
Population aged 15-49 (thousands)	2004	42,423	UN population division database
Annual population growth rate (%)	1992-2002	2.1	UN population division database
% of population in urban areas	2003	60.6	UN population division database
Average annual growth rate of urban population	2000-2005	3.1	UN population division database
Crude birth rate (births per 1,000 pop.)	2004	24.5	UN population division database
Crude death rate (deaths per 1,000 pop.)	2004	5.1	UN population division database
Maternal mortality rate (per 100,000 live births)	2000	200	WHO (WHR2004)/UNICEF
Life expectancy at birth (years)	2002	68.3	World Health Report 2004, WHO
Total fertility rate	2002	3.2	World Health Report 2004, WHO
Infant mortality rate (per 1,000 live births)	2000	28	World Health Report 2004, WHO
Under 5 mortality rate (per 1,000 live births)	2000	40	World Health Report 2004, WHO

SOCIO-ECONOMIC DATA	YEAR	ESTIMATE	SOURCE
Gross national income, ppp, per capita (Int.\$)	2002	4,280	World Bank
Gross domestic product, per capita % growth	2001-2002	2.4	World Bank
Per capita total expenditure on health (Int.\$)	2001	169	World Health Report 2004, WHO
General government expenditure on health as % of total expenditure on health	2001	45.2	World Health Report 2004, WHO
Total adult illiteracy rate	2000	5.1	UNESCO
Adult male illiteracy rate	2000	4.9	UNESCO
Adult female illiteracy rate	2000	5.2	UNESCO
Gross primary school enrolment ratio, male	2000/2001	113	UNESCO
Gross primary school enrolment ratio, female	2000/2001	113	UNESCO
Gross secondary school enrolment ratio, male	2000/2001	74	UNESCO
Gross secondary school enrolment ratio, female	2000/2001	81	UNESCO

### Contact address

UNAIDS/WHO Working Group on Global  
HIV/AIDS and STI Surveillance  
20, Avenue Appia  
CH - 1211 Geneva 27  
Switzerland  
Fax: +41-22-791-4834

email: [hivstrategicinfo@who.int](mailto:hivstrategicinfo@who.int) or  
[estimates@unaids.org](mailto:estimates@unaids.org)

website: <http://www.who.int/hiv>

<http://www.unaids.org>

Extracts of the information contained in these fact sheets may be reviewed, reproduced or translated for research or private study but not for sale or for use in conjunction with commercial purposes. Any use of information in these fact sheets should be accompanied by the following acknowledgment "UNAIDS/WHO epidemiological fact sheets on HIV/AIDS and Sexually Transmitted Infections, 2004 Update".

## HIV prevalence in different populations

This section contains information about HIV prevalence in different populations. The data reported in the tables below are mainly based on the HIV database maintained by the United States Bureau of the Census where data from different sources, including national reports, scientific publications and international conferences are compiled. To provide a simple overview of the current situation and trends over time, summary data are given by population group, geographical area (Major Urban Areas versus Outside Major Urban Areas), and year of survey. Studies conducted in the same year are aggregated and the median prevalence rates (in percentages) are given for each of the categories. The maximum and minimum prevalence rates observed, as well as the total number of surveys/sentinel sites, are provided with the median, to give an overview of the diversity of HIV-prevalence results in a given population within the country. Data by sentinel site or specific study from which the medians were calculated are printed at the end of this fact sheet.

The differentiation between the two geographical areas Major Urban Areas and Outside Major Urban Areas is not based on strict criteria, such as the number of inhabitants. For most countries, Major Urban Areas were considered to be the capital city and - where applicable - other metropolitan areas with similar socio-economic patterns. The term Outside Major Urban Areas considers that most sentinel sites are not located in strictly rural areas, even if they are located in somewhat rural districts.

### HIV sentinel surveillance\*


Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		
Pregnant women																					
Sex workers	Major urban areas	N-Sites	2.00					1.00	2.00	1.00								2.00			
		Minimum	0					0.06	0.33	0.33									0		
		Median	0.02					0.06	0.44	0.33									0		
		Maximum	0.04					0.06	0.54	0.33									0		
	Outside major urban areas	N-Sites	15.00	11.00	15.00				1.00	2.00			1.00	1.00	1.00	1.00	2.00				
		Minimum	0	0	0				0	0			0	0	0	0	0				
		Median	0	0	0				0	0			0	0	0	0	0				
		Maximum	0.48	0.68	0.02				0	0			0	0	0	0	0				
Injecting drug users	Major urban areas	N-Sites							1.00	1.00											
		Minimum							0	0											
		Median							0	0											
		Maximum							0	0											
	Outside major urban areas	N-Sites						1.00	1.00	2.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00				
		Minimum						0	0	0	0	0.30	0	0	0	0	0				
		Median						0	0	0	0	0.30	0	0	0	0	0				
		Maximum						0	0	0	0	0.30	0	0	0	0	0				
STI patients	Major urban areas	N-Sites							1.00	1.00											
		Minimum							0	0											
		Median							0	0											
		Maximum							0	0											
	Outside major urban areas	N-Sites						1.00	1.00	1.00											
		Minimum						0	0	0											
		Median						0	0	0											
		Maximum						0	0	0											
Men having sex with men	Major urban areas	N-Sites							1.00	1.00											
		Minimum							0	0											
		Median							0	0											
		Maximum							0	0											
	Outside major urban areas	N-Sites	2.00						1.00	3.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00				
		Minimum	0						0	0	0.20	0	0.29	0	0	0	0				
		Median	0.06						0	0	0.20	0	0.29	0	0	0	0				
		Maximum	0.12						0	0	0.20	0	0.29	0	0	0	0				
Tuberculosis patients	Major urban areas	N-Sites								1.00											
		Minimum								2.16											
		Median								2.16											
		Maximum								2.16											

\*Detailed data by site can be found in the Annex.

## Maps & charts

Mapping the geographical distribution of HIV prevalence among different population groups may assist in interpreting both the national coverage of the HIV surveillance system as well in explaining differences in levels of prevalence. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, in collaboration with the WHO Public Health Mapping Team, Communicable Diseases, is producing maps showing the location and HIV prevalence in relation to population density, major urban areas and communication routes. For generalized epidemics, these maps show the location of prevalence of antenatal surveillance sites.

Trends in antenatal sentinel surveillance for higher prevalence countries, or in prevalence among selected populations for countries with concentrated epidemics, are a new addition. These are presented for those countries where sufficient data exist.



## Reported AIDS cases

Following WHO and UNAIDS recommendations, AIDS case reporting is carried out in most countries. Data from individual AIDS cases are aggregated at the national level and sent to WHO. However, case reports come from surveillance systems of varying quality. Reporting rates vary substantially from country to country and low reporting rates are common in developing countries due to weaknesses in the health care and epidemiological systems. In addition, countries use different AIDS case definitions. A main disadvantage of AIDS case reporting is that it only provides information on transmission patterns and levels of infection approximately 5-10 years in the past, limiting its usefulness for monitoring recent HIV infections.

Despite these caveats, AIDS case reporting remains an important advocacy tool and is useful in estimating the burden of HIV-related morbidity as well as for short-term planning of health care services. AIDS case reports also provide information on the demographic and geographic characteristics of the affected population and on the relative importance of the various exposure risks. In some situations, AIDS reports can be used to estimate earlier HIV infection patterns using back-calculation. AIDS case reports and AIDS deaths have been dramatically reduced in industrialized countries with the introduction of Anti-Retroviral Therapy (ART).

1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998
					2	4	11	13	11	10	18	17	21	38	57	51	50	23	45
1999	2000	2001	2002	2003	Total	UNK	Date of last report												
78	40	56	44	47	636		12/31/2003												

## Curable sexually transmitted infections (STIs)

The predominant mode of transmission of both HIV and other STIs is sexual intercourse. Measures for preventing sexual transmission of HIV and STIs are the same, as are the target audiences for interventions. In addition, strong evidence supports several biological mechanisms through which STIs facilitate HIV transmission by increasing both HIV infectiousness and HIV susceptibility. Thus, detection and treatment of individuals with STIs is an important part of an HIV control strategy. In summary, if the incidence/prevalence of STIs is high in a country, then there is the possibility of high rates of sexual transmission of HIV. Monitoring trends in STIs provides valuable insight into the likelihood of the importance of sexual transmission of HIV within a country, and is part of second generation surveillance. These trends also assist in assessing the impact of behavioural interventions, such as delaying sexual debut, reducing the number of sex partners and promoting condom use.

Clinical services offering STI care are an important access point for people at high risk for both STIs and HIV. Identifying people with STIs allows for not only the benefit of treating the STI, but for prevention education, HIV testing, identifying HIV-infected persons in need of care, and partner notification for STIs or HIV infection. Consequently, monitoring different components of STI prevention and control can also provide information on HIV prevention and control activities within a country.

### STI syndromes

Reported cases	1996	1997	1998	1999	2000	2001	2002	2003	Incidence 2003
----------------	------	------	------	------	------	------	------	------	----------------

Comments:

Source:

### Syphilis prevalence, women

Percent of blood samples taken from pregnant women aged 15-49 that test positive for syphilis - positive reaginic and treponemal test-during routine screening at selected antenatal clinics.

Year	Area	Rate	Range
------	------	------	-------

Comments:

Source:

### Estimated prevalence of curable STIs among female sex workers

#### - Chlamydia

Year	Area	Rate	Range
1997-1999	Urban/Rural	35	27-35

Comments:

Source: Department of Health Philippines, WHO Regional Office for the Western Pacific. Consensus Report on STI, HIV and AIDS Epidemiology 2000.

#### - Gonorrhoea

Year	Area	Rate	Range
1997-1999	Urban/Rural	10.95	2.6-38

Comments:

Source: Department of Health Philippines, WHO Regional Office for the Western Pacific. Consensus Report on STI, HIV and AIDS Epidemiology 2000.

Estimated prevalence of curable STIs among female sex workers (continued)**- Syphilis**

Year	Area	Rate	Range
------	------	------	-------

**Comments:**

**Source:**

**- Trichomoniasis**

Year	Area	Rate	Range
1997-1999	Urban/Rural	6.5	5-8

**Comments:**

**Source:** Department of Health Philippines, WHO Regional Office for the Western Pacific. Consensus Report on STI, HIV and AIDS Epidemiology 2000.

## Health service and care indicators

HIV prevention strategies depend on the twin efforts of care and support for those living with HIV or AIDS, and targeted prevention for all people at risk or vulnerable to the infection. It is difficult to capture such a large range of activities with one or just a few indicators. However, a set of well-established health care indicators may help to identify general strengths and weaknesses of health systems. Specific indicators, such as access to testing and blood screening for HIV, help to measure the capacity of health services to respond to HIV/AIDS - related issues.

### Access to health care

Indicators	Year	Estimate	Source
% of population with access to health services - total			
% of population with access to health services - urban			
% of population with access to health services - rural			
Contraceptive prevalence rate (%)	1998	46	UNICEF/UNPOP
Percentage of contraceptive users using condoms			
% of births attended by skilled health personnel	2000	56.4	WHO
% of 1-yr-old children fully immunized - DPT	2002	70	WHO/UNICEF
% of 1-yr-old children fully immunized - Measles	2001	75	WHO/UNICEF
% of ANC clinics where HIV testing is available			

### Number of adults (15-49) with advanced HIV infection receiving ARV therapy as of June 2004

#### Adults on treatment

Number:	71
Source:	WHO

### Estimated number of adults (15-49) in need of treatment in 2003

#### Adults needing treatment

Number:	710
Source:	WHO/UNAIDS

### Coverage of HIV testing and counselling

Number of public and NGO services providing testing and counselling services.

Year	Area	N=
------	------	----

Comments:

Source:

## Knowledge and behaviour

In most countries the HIV epidemic is driven by behaviours (e.g.: multiple sexual partners, injecting drug use) that expose individuals to the risk of infection. Information on knowledge and on the level and intensity of risk behaviour related to HIV/AIDS is essential in identifying populations most at risk for HIV infection and in better understanding the dynamics of the epidemic. It is also critical information in assessing changes over time as a result of prevention efforts. One of the main goals of the 2nd generation HIV surveillance systems is the promotion of a standard set of indicators defined in the National Guide (Source: National AIDS Programmes, A Guide to Monitoring and Evaluation, UNAIDS/00.17) and regular behavioural surveys in order to monitor trends in behaviours and to target interventions.

The indicators on knowledge and misconceptions are an important prerequisite for prevention programmes to focus on increasing people's knowledge about sexual transmission, and, to overcome the misconceptions that act as a disincentive to behaviour change. Indicators on sexual behaviour and the promotion of safer sexual behaviour are at the core of AIDS programmes, particularly with young people who are not yet sexually active or are embarking on their sexual lives, and who are more amenable to behavioural change than adults. Finally, higher risk male-male sex reports on unprotected anal intercourse, the highest risk behaviour for HIV among men who have sex with men.

### Knowledge of HIV prevention methods

Prevention indicator: Percentage of young people 15-24 who both correctly identify two ways of preventing the sexual transmission of HIV and who reject three misconceptions about HIV transmission.

Year	Male	Female

Comments:

Source:

### Reported condom use at last higher risk sex (young people 15-24)

Prevention indicator: Proportion of young people reporting the use of a condom during sex with a non-regular partner.

Year	Male	Female

Comments: For this indicator only data will be shown if they were collected after 1998.

Source:

### Age-mixing in sexual partnerships among young women

The proportion of young women who have had sex in the last 12 months with a partner who is 10 or more years older than themselves.

Year	Area	Age group	Male	Female	All

Comments:

Source:

### Reported non-regular sexual partnerships

Prevention indicator: Proportion of young people 15-24 having at least one sex partner other than a regular partner in the last 12 months.

Year	Male	Female

Comments:

Source:

Knowledge and behaviour (continued)Ever used a condom

Percentage of people who ever used a condom.

Year	Area	Age group	Male	Female	All
------	------	-----------	------	--------	-----

Comments:

Source:

Adolescent pregnancy

Percentage of teenagers 15-19 who are mothers or pregnant with their first child.

Year	Percentage
------	------------

Comments:

Source:

Age at first sexual experience

Proportion of 15-19 year olds who have had sex before age 15.

Year	Male	Female
1998		1

Comments:

Source: DHS

## Prevention indicators

Male and female condoms are the only technology available that can prevent sexual transmission of HIV and other STIs. Persons exposing themselves to the risk of sexual transmission of HIV should have consistent access to high quality condoms. AIDS Programs implement activities to increase both availability of and access to condoms. These activities should be monitored and have resources directed to problem areas. The indicator below highlights the availability of condoms. However, even if condoms are widely available, this does not mean that individuals can or do access them.

### Condom availability nationwide

Total number of condoms available for distribution nationwide during the preceding 12 months, divided by the total population aged 15-49.

Year	N	Rate
------	---	------

Comments:

Source:

### Prevention of mother-to-child transmission (MTCT) nationwide

Percentage of women who were counselled during antenatal care for their most recent pregnancy, accepted an offer of testing and received their test results, of all women who were pregnant at any time in the preceding two years.

Year	N	Rate
------	---	------

Comments:

Source:

Blood safety programs aim to ensure that the majority of blood units are screened for HIV and other infectious agents. This indicator gives an idea of the overall percentage of blood units that have been screened to high enough standards that they can confidently be declared free of HIV.

### Screening of blood transfusions nationwide

Percentage of blood units transfused in the last 12 months that have been adequately screened for HIV according to national or WHO guidelines.

Year	N	Rate
------	---	------

Comments:

Source:

## Sources

Data presented in this Epidemiological Fact Sheet come from several sources, including global, regional and country reports, published documents and articles, posters and presentations at international conferences, and estimates produced by UNAIDS, WHO and other United Nations agencies. This section contains a list of the more relevant sources used for the preparation of the Fact Sheet. Where available, it also lists selected national Web sites where additional information on HIV/AIDS and STI are presented and regularly updated. However, UNAIDS and WHO do not warrant that the information in these sites is complete and correct and shall not be liable whatsoever for any damages incurred as a result of their use.

- Basaca-Sevilla, V., C. G. Hayes, G. Espinosa, et al. 1989 The Search for the Prevalence of Infection with the Human Immunodeficiency Virus, The Hepatitis B Virus and Treponema Pallidum.. *Journal of Philippine Medical Association*, vol. 65, no. 2, pp. 223-234.
- Badoy, T. J., P. Ramos-Jimenez, A. Panadero 1997 Tracking the HIV Epidemic in the Philippines - Methods, Relevant Findings, and Directions in HIV Prevention and Control Program Paper Presented at the UNAIDS Regional Workshop on "Evidence of Behavioural Change in the Context of HIV Decline in Thailand," 19-22 May, Bangkok, Thailand.
- Cueto, R. C. 2001 Syphilis Prevalence among Commercial Sex Workers in Davao City, Philippines 6th International Congress on AIDS in Asia and the Pacific, Melbourne, Australia, 10/5-10, Abstract Mo0997.
- Carabana, C. L. H., R. Silapan, R. Mendoza 2001 5th HIV/AIDS Serologic Surveillance among Female Sex Workers in Zamboanga City, Philippines 6th International Congress on AIDS in Asia and the Pacific, Melbourne, Australia, 10/5-10, Poster Mo1197.
- Dayrit, M. M., O. T. Monzon., V. Basaca-Sevilla, et al. 1987 Emerging Patterns of HIV Infection and Control in the Philippines *Western Journal of Medicine*, vol. 147, pp. 723-725.
- Dayrit, M. M., T. J. Badoy 1994 Results of the Second Round of Surveillance Activities, NCR-Quezon, City, Metro Cebu, and Metro Davao, April - June, 1994 National HIV Sentinel Surveillance System, Department of Health.
- Dayrit, M. M., M. E. White, T. J. Badoy 1993 Results of the First HIV Surveillance Round National Capital Region and Metro Cebu, June - September 1993 National HIV Sentinel Surveillance System, Department of Health.
- Harare, H. 1987 AIDS Powderkeg Waiting to Blow Up Across Asia *Harare Herald*, July 27, p. 8.
- Reyes, C. C., E. N. Fabro, N. M. Mariano, et al. 1995 Preliminary Study on the Incidence of HIV Seropositivity amongst High-Risk Filipino PTB Patients IUVDT World STD/AIDS Congress, Singapore, 3/19-23, Free Paper 9.
- San Juan, A., R. Mangonon, P. Perez 2002 Low and Slow Transmission of HIV in Pasay City, Philippines XIV International AIDS Conference, Barcelona, Spain, 7/7-12, Abstract MoPeC3506.
- Tan, M. L., M. M. Dayrit 1994 HIV/AIDS in the Philippines *AIDS*, vol. 8, suppl. 2, pp. S125-S130.
- World Health Organization, Western Pacific Region 1993 AIDS Surveillance Report Global Programme on AIDS, vol. 1, no. 1, July, p. 2.
- World Health Organization 2000 HIV and Behavioural Surveillance in the Philippines STI/HIV/AIDS Surveillance Report, no. 15, pp. 5-6.
- Yeoh, E. 1990 The Growing Problem of AIDS in Asia VI International Conference on AIDS, San Francisco, 6/24, Closing Ceremony, vol. 3, p. 93.

### **Websites:**

## Annex: HIV surveillance by site

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003		
Pregnant women																				
Sex workers	Major urban areas	Metro Manila	0.04																	
		National						0.06												
		Pasay							0.54									0		
			Quezon City	0					0.33	0.33										
	Outside major urban areas		Alaminos, Pangasinan	0																
			Angeles/Mabalacat	0.03	0.03	0														
			Bacolod			0														
			Baguio, Baguio City	0	0	0														
			Batangas	0		0														
			Bocaue	0	0	0														
			Cagayan de Oro																	
			Caloocan																	
			Cavite	0		0														
			Cebu	0	0															
			Dagupan, Pangasinan	0		0														
			Davao																	
			Davao area		0															
			Davao City																0	
			General Santos																	
			Gumaca, Lucena			0														
			Iloilo, Iloilo		0															
			Kalibo/Aklan, Kalibo, Aklan																	
			Lapu-Lapu																	
			Legaspi																	
			Los Banos, Laguna	0		0														
			Lucena	0		0														
			Makati	0																
			Marikina	0																
			Metro Cebu							0	0									
		Metro Davao								0										
		Naga																		
		Olongapo/Subic	0.07	0.06	0.02															
	Roxas																			
	San Antonio		0	0																
	San Fernando	0.48	0.65	0																
	Tacloban																			
	Tuguegarao			0																
	Urdaneta, Pangasinan	0		0																
	Zamboanga		0								0	0	0	0	0					
Injecting drug users	Major urban areas	Quezon City						0	0											
	Outside major urban areas	Baguio & Cebu					0													
		Metro Cebu							0	0										
		Not specified (1), Not specified						0	0	0.30	0	0	0	0	0					
STI patients	Major urban areas	Quezon City						0	0											
	Outside major urban areas	Baguio & Cebu					0													
		Metro Cebu						0	0											
Men having sex with men	Major urban areas	Quezon City						0	0											
	Outside major urban areas	Metro Cebu						0	0											
		Metro Davao							0	0										
		Not specified (1), Not specified	0.12						0	0.20	0	0.29	0	0	0					
	Not specified (2), Not specified	0																		
Tuberculosis patients	Major urban areas	Manila						2.16												